

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS**



All Weather
PCT
First Formula*
BLOCKS RODENTICIDE



ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting

CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 150g



HOLDINGS PTY LTD

APVMA Approval No: 59875/150/0607

***PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD
(PCT Holdings Pty Ltd ABN 11 099 023 962)**

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

DIRECTIONS FOR USE

RESTRAINTS

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait in exposed areas.

DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence, whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/150/0607

Batch No.

D.O.M



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS



All Weather
PCT
First Formula*
BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting

CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 300g



APVMA Approval No: 59875/300/0607

***PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD
(PCT Holdings Pty Ltd ABN 11 099 023 962)**

**1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>
CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877**

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait in exposed areas.

DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/300/0607

Batch No.

D.O.M



POISON**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS****All Weather
PCT
First Formula*
BLOCKS RODENTICIDE****ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)**

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

**TO CONTROL RATS AND MICE
AS PER THE DIRECTIONS FOR USE TABLE****One Feed – Fast Acting**

CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 1kg

APVMA Approval No: 59875/1/0607

*PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD
(PCT Holdings Pty Ltd ABN 11 099 023 962)1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE
RESTRAINTS**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.
DO NOT place bait in exposed areas.
DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.****GENERAL INSTRUCTIONS**

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.**PRECAUTION:**

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/1/0607
Batch No. D.O.M.



POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS**

All Weather
PCT First Formula*
BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting

CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 2.4kg



APVMA Approval No: 59875/2.4/0607

*PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD (PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

First Formula 2.4kg 116 mm x 180 mm

DIRECTIONS FOR USE RESTRAINTS

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait in exposed areas.

DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/2.4/0607

Batch No.

D.O.M.





POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS**

All Weather
PCT First Formula*
BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting

CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 5kg



HOLDINGS PTY LTD

APVMA Approval No: 59875/5/0607

*PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD (PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

First Formula 5kg 116 mm x 180 mm

DIRECTIONS FOR USE RESTRAINTS

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.
DO NOT place bait in exposed areas.
DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/5/0607

Batch No.

D.O.M.



POISON**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS****All Weather**

PCT First Formula*

BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting
 CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 8kg

PCT
 HOLDINGS PTY LTD

APVMA Approval No: 59875/8/0607

*PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD
 (PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>
 CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE
RESTRAINTS**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.
 DO NOT place bait in exposed areas.
 DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/8/0607

Batch No. D.O.M.

POISON**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS**

All Weather
PCT
First Formula*
BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings, exterior and interior of buildings and structures.

TO CONTROL RATS AND MICE AS PER THE DIRECTIONS FOR USE TABLE

One Feed – Fast Acting
CONTROLS SPECIES RESISTANT TO WARFARIN

Net Contents 10kg

PCT
HOLDINGS PTY LTD

APVMA Approval No: 59875/10/0607

*PCT First Formula is a registered trademark of PCT INTERNATIONAL PTY LTD
(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE
RESTRAINTS**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.
DO NOT place bait in exposed areas.
DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial buildings.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1-3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

PCT First Formula Blocks can be used both indoors and outdoors. It is recommended that clearly marked tamper evident stations be used wherever baits are to be placed. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Look for dead rats or mice – burn or bury them, do not place in refuse bins. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTION:

Avoid contact with food, food utensils, or places where food is prepared or stored.

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. Place baits in locations inaccessible to other animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Place baits in locations inaccessible to animals other than target animals. The product is restricted to use in and around buildings (within 2 metres), structures or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If not available bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers that have held bait for any other purpose.

Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS: Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Vitamin K¹ (PHYTOMENADIONE) is antidotal.

MSDS: For further information please refer to the Material Safety Data Sheet.

Contains human taste deterrent that deters accidental consumption by children.

CONDITIONS OF SALE

The use of All Weather PCT First Formula Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT INTERNATIONAL PTY LTD regarding its suitability or efficacy for any purpose. PCT INTERNATIONAL cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 59875/10/0607

Batch No.

D.O.M.