Australian Pesticides & Veterinary Medicines Authority

Product Name: Gnaw Rodenticide
APVMA Approval No: 94661/151609

Label Name:	Gnaw Rodenticide

Signal Headings:	DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	ACTIVE CONSTITUENT: 0.75 g/kg CHOLECALCIFEROL Contains DENATONIUM BENZOATE (BITTERING AGENT)

Mode of Action:	

Statement of Claims:	For the control of rats and mice, including anticoagulant resistant species, in and around domestic, commercial and agricultural buildings and for the control of rats within macadamia orchards.

Net Contents:	150 g - 20 kg

Restraints:	RESTRAINTS: DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT place bait stations in open and bait must only be used in lockable, tamper-resistant bait stations. DO NOT use this product for the control of protected native rodents, or other native animals. DO NOT apply within native vegetation.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	

Trade Advice:	

General Instructions:	GENERAL INSTRUCTIONS Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. In Crop Situations Baiting within macadamia orchards should be part of a comprehensive Integrated Pest Management (IPM) strategy, which focusses on the elimination of alternative food resources, the encouragement of natural predators, weed control strategies (around and within crop), effective rodent monitoring, clearance or revegetation of adjacent scrub land and strategically timed baiting. Pre-baiting strategies are not required when using Gnaw Rodenticide.

Resistance Warning:	

Precautions:	PRECAUTIONS This product is only to be utilised by placement in lockable bait stations. DO NOT place baits in areas accessible to children or pets. DO NOT place in positions where there is a possibility of contaminating food or on surfaces that come into direct contact with food.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. DO NOT place baits in locations that are accessible to domestic animals, livestock, non-target native animals or birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

Safety Directions:	SAFETY DIRECTIONS Wash hands after use.

First Aid Instructions:	FIRST AID If poisoning occurs get to a doctor or hospital quickly.

First Aid Warnings:	

DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, hospitals, inside transport vehicles (including ships) around grain terminals, storage areas and perimeter fence lines. At food facilities, for fence line use only.	RATS *(Rattus norvegicus, Rattus rattus)*	3 - 9 blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Place baits in lockable bait stations. Inspect bait stations on a regular basis. Replace eaten baits. If eaten quickly increase number of baits. Continue observation and replenishment until no more baits are taken.
	MICE *(Mus musculus)*	1 - 2 blocks per bait station.	
Macadamia orchards	RATS *(Rattus rattus* also known as Black rat, Roof rat, Ship rat or Fruit rat)	3 - 4 blocks per bait station.	Baiting programs within orchards should be part of a comprehensive IPM program. Headlands and adjacent non-crop habitats should be kept weed free and slashed (where applicable) to expose rodents to natural predators. Monitor rodent activity and damage levels prior to baiting. Concentrate baiting efforts to the outer rows of orchard blocks adjacent to non-crop habitats or in areas of the orchard with signs of high activity. Place baits in bait stations in areas where rodents frequent, including in trees, on the orchard ground layer and within adjacent non-crop habitats. Bait stations should be placed between 8-12 m apart in areas where rodent activity has been observed. Ensure bait stations are placed to minimise the ingress of water such that baits remain dry. Check baits every 7-14 days and once feeding has stopped baits may be removed. Remove baits if there is evidence of native wildlife visiting bait stations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.