


# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

NET CONTENTS 150g



**PCT**  
HOLDINGS PTY LTD

APVMA Approval No: 64849/150g/0410

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD

(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait stations in open.

SITUATION	PEST	RATE	Critical Comments
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals).	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS**

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

**PRECAUTION: KEEP OUT OF REACH OF CHILDREN**

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

**STORAGE AND DISPOSAL**

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

**MSDS**

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

**APVMA Approval No. 64849/150g/0410**

**B/N**

**DOM**

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## BROMA BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

**NET CONTENTS 300g**



**APVMA Approval No: 64849/300g/0410**

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD

(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait stations in open.

SITUATION	PEST	RATE	Critical Comments
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals),	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS**

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

**PRECAUTION: KEEP OUT OF REACH OF CHILDREN**

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

**STORAGE AND DISPOSAL**

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

**MSDS**

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.


**CONDITIONS OF SALE**

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

**APVMA Approval No. 64849/300g/0410**


**B/N**

**DOM**



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

**NET CONTENTS 1 kg**



APVMA Approval No: 64849/1kg/0410

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD

(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.  
DO NOT place bait stations in open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals).	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS**

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

**PRECAUTION: KEEP OUT OF REACH OF CHILDREN**

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

**STORAGE AND DISPOSAL**

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidental.

**MSDS**

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

**APVMA Approval No. 64849/1kg/0410**

**B/N DOM**



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## BROMA BLOCKS RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

**NET CONTENTS 2.4 kg**



APVMA Approval No: 64849/2.4kg/0410

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD  
(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.  
DO NOT place bait stations in open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals).	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS**

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

**PRECAUTION: KEEP OUT OF REACH OF CHILDREN**

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

**STORAGE AND DISPOSAL**

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

**MSDS**

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

**APVMA Approval No. 64849/2.4kg/0410**

**B/N**

**DOM**



# SureFire<sup>®</sup>

## BROMA BLOCKS RODENTICIDE

NET CONTENTS 8kg



PCT  
HOLDINGS PTY LTD

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

NET CONTENTS 8 kg

  
**PCT**  
HOLDINGS PTY LTD

APVMA Approval No: 64849/8kg/0210

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD  
(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877



## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait stations in open.

SITUATION	PEST	RATE	Critical Comments
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals).	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

### GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

### PRECAUTION: KEEP OUT OF REACH OF CHILDREN

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

## PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

### STORAGE AND DISPOSAL

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

### MSDS

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

### CONDITIONS OF SALE

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 64849/8kg/0410

B/N

DOM



# SureFire<sup>®</sup>

## BROMA BLOCKS RODENTICIDE

NET CONTENTS 10kg



PCT  
HOLDINGS PTY LTD

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE  
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice, including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

NET CONTENTS 10 kg



APVMA Approval No: 64849/10kg/0210

\*SUREFIRE® BROMA BLOCKS is a registered trademark of PCT INTERNATIONAL PTY LTD  
(PCT Holdings Pty Ltd ABN 11 099 023 962)

1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877



## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait stations in open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships and around grain terminals).	RATS ( <i>Rattus norvegicus</i> , <i>Rattus rattus</i> )	4 or more blocks per bait station.	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks is necessary to reduce rat/mouse numbers.
	MICE ( <i>Mus musculus</i> )	2 to 4 blocks per infested area.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

### GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

### PRECAUTION: KEEP OUT OF REACH OF CHILDREN

DO NOT allow bait to contaminate foodstuffs or feed intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.

The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

## PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

### STORAGE AND DISPOSAL

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS:** Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

### MSDS

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

### CONDITIONS OF SALE

The use of Surefire Broma Blocks Rodenticide is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by PCT HOLDINGS PTY LTD regarding its suitability or efficacy for any purpose. PCT HOLDINGS PTY LTD cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No. 64849/10kg/0410

B/N

DOM